arXiv:1412.8194v3 [math.AT] 2 Oct 2015

HOMOLOGY GROUPS OF SPACES OF NON-RESULTANT QUADRATIC POLYNOMIAL SYSTEMS IN $\mathbb{R}^3$

V.A. VASSILIEV

Abstract. We calculate rational homology groups of spaces of non-resultant (i.e. having no non-trivial common zeros) systems of homogeneous quadratic polynomials in $\mathbb{R}^3$.

1. Introduction

The *resultant subvariety* in the space of systems of homogeneous polynomials in $\mathbb{R}^n$ consists of systems having common zeros besides the origin in $\mathbb{R}^n$. The varieties of this kind were considered, in particular, by S. Smale and his collaborators in connection with the $P \stackrel{?}{=} NP$ problem over fields of zero characteristic, see [13], [4], [12]. Also, the topology of complements of resultant varieties in certain spaces of systems of non-homogeneous polynomials in one complex variable was studied in [7], [9], [17]. The homology groups of spaces of non-resultant homogeneous systems in $\mathbb{R}^2$ were calculated in [21]. Below we study the next complicated problem — of homogeneous quadratic polynomial systems in $\mathbb{R}^3$.

Let $W \simeq \mathbb{R}^6$ be the space of all real quadratic forms in $\mathbb{R}^3$. For any natural k, consider the space $W^k \simeq \mathbb{R}^{6k}$ of all quadratic maps $\mathbb{R}^3 \to \mathbb{R}^k$. Denote by Σ the *resultant subvariety* in W^k, i.e. the space of all systems of k quadratic forms having common zeros in $\mathbb{R}^3 \setminus 0$.

Theorem 1. *For any even $k \geq 4$, the Poincaré polynomial of the group $H^*(W^k \setminus \Sigma, \mathbb{Q})$ is equal to*

$$(1 + t^{k-1})(1 + t^{3k-9} + t^{5k-14}). \tag{1}$$

For any odd $k \geq 3$, the Poincaré polynomial of $H^*(W^k \setminus \Sigma, \mathbb{Q})$ is equal to

$$1 + t^{k-1} + t^{3k-7}(1 + t^{k-5})(1 + t^{2k-3}). \tag{2}$$

The group $H^i(W^2 \setminus \Sigma, \mathbb{Q})$ is equal to $\mathbb{Q}$ if $i = 0$ or $i = 1$, and is trivial otherwise.

Supported by RSF grant 14-50-00005.

Example 1. For $k = 3$, the polynomial (2) is equal to $(2 + t^3)(1 + t^2)$. In particular, the space $W^3 \setminus \Sigma$ consists of two connected components. Indeed, any non-resultant triple of quadratic forms in $\mathbb{R}^3$ defines a map $\mathbb{RP}^2 \to \mathbb{R}^3 \setminus 0 \sim S^2$. The mod 2 degree of this map is invariant along any connected component of $W^3 \setminus \Sigma$. This degree is equal to 0 for any system containing non-negative quadratic forms, and is non-trivial for a triple of quadratic forms whose zero sets in $\mathbb{RP}^2$ meet like .

2. First reductions

2.1. **Alexander duality.** Denote by $\overline{H}_*$ the *Borel–Moore homology group*, i.e. the homology group of the complex of locally finite singular chains of a topological space. In the case of homology groups with coefficients in a group (i.e. in a constant local system) A, an equivalent definition is as follows: $\overline{H}_*(X, A)$ is the homology group of the one-point compactification of X reduced modulo the added point. However, in the case of non-constant coefficient systems the latter definition is generally not applicable. $\tilde{H}_*$ and $\tilde{H}^*$ is the notation for the homology and cohomology groups reduced modulo a point.

Following [1], we use the Alexander isomorphism

$$\tilde{H}^i(W^k \setminus \Sigma) \simeq \overline{H}_{6k-i-1}(\Sigma). \tag{3}$$

2.2. **A toy calculation: homology of Stiefel manifolds.** To calculate the right-hand group in (3) we consider a *simplicial resolution* of the space Σ. Let us demonstrate this method in a simpler situation, replacing quadratic maps by linear ones. The space of non-resultant collections of k linear functions $\mathbb{R}^3 \to \mathbb{R}^1$, $k \geq 3$, is homotopy equivalent to the Stiefel manifold $V_3(\mathbb{R}^k)$. Its cohomology group is well-known; let us show how to calculate it by our techniques.

Proposition 1. *For any $k \geq 3$, the Poincaré polynomial of the group $H^i(V_3(\mathbb{R}^k), \mathbb{Q})$ is equal to $(1 + t^{k-1})(1 + t^{2k-5})$ if k is even, and to $(1 + t^{k-3})(1 + t^{2k-3})$ if k is odd.*

Proof. Consider the entire space $\mathbb{R}^{3k}$ of collections of k linear functions, and the discriminant variety Σ in it consisting of forms vanishing at some non-trivial subspace. To construct the simplicial resolution σ of this variety, we list all subsets in $\mathbb{RP}^2$ which can be distinguished by systems of linear equations.

It are 1) particular points (the space of them is $\mathbb{RP}^2$ itself), 2) the lines (the space of them is the dual projective plane $\widehat{\mathbb{RP}^2}$), and 3) entire

$\mathbb{RP}^2$: this is a single object named $[\mathbb{RP}^2]$. Let us denote these spaces of subsets by J_1, J_2 and J_3 respectively.

We consider them as topological spaces with their standard topology. The disjoint union $J_1 \sqcup J_2 \sqcup J_3$ is a partially ordered set with the order defined by inclusion of corresponding subsets. Consider the topological order complex Ψ of this poset. Namely, take the join $J_1 * J_2 * J_3$ of these three spaces (i.e., the naturally topologized union of triangles and segments, whose vertices correspond to the points of different spaces) and define the space Ψ as the union of only those of these simplices, all whose vertices are incident to one another. For any point X of J_1, J_2 or J_3, define its subordinate order subcomplex $\Psi(X)$ as the union of all such simplices, all whose vertices correspond to sets contained in the set $\{X\}$. Also, denote by $\partial\Psi(X)$ the *link* of $\Psi(X)$, i.e. the union of all simplices participating in $\Psi(X)$ but not containing its vertex X. Let $\breve{\Psi}(X)$ be the difference $\Psi(X) \setminus \partial\Psi(X)$. Denote by Ψ_p, $p = 1, 2, 3$ the union of all complexes $\Psi(X)$ over all $X \in J_j$, $j \leq p$. Obviously, $\Psi_3 = \Psi([\mathbb{RP}^2]) = \Psi$.

The *simplicial resolution* σ of Σ is defined as a subspace in the direct product $\Psi \times \mathbb{R}^{3k}$. Namely, for any $p = 1, 2$, or 3, and any $X \in J_p$ we take the product $\Psi(X) \times L(X)$, where $L(X) \subset \mathbb{R}^{3k}$ is the space of all systems of k linear functions vanishing on X. σ is defined as the union of all such products over all p and all $X \in J_p$. The subspace $\sigma_p \subset \sigma$ is the union of such products only over all $X \in J_j$, $j \leq p$. It is easy to see that the difference $\sigma_p \setminus \sigma_{p-1}$ is the space of a fiber bundle over J_p; its fiber over $X \in J_p$ is equal to $\breve{\Psi}(X) \times L(X)$. In particular, $\sigma_p \setminus \sigma_{p-1}$ is the space of a vector bundle over $\Psi_p \setminus \Psi_{p-1}$ with the fibers equal to $L(X)$, $X \in J_p$.

The projection $\sigma \to \Sigma$ is proper with contractible fibers, hence defines an isomorphism $\overline{H}_*(\sigma, \mathbb{Q}) \to \overline{H}_*(\Sigma, \mathbb{Q})$. Now let us calculate the spectral sequence converging to the group $\overline{H}_*(\sigma, \mathbb{Q})$ and generated by the filtration $\sigma_1 \subset \sigma_2 \subset \sigma_3 \equiv \sigma$. By definition, its initial term $E^1_{p,q}$ is equal to $\overline{H}_{p+q}(\sigma_p \setminus \sigma_{p-1}, \mathbb{Q})$.

The space σ_1 is fibered over $J_1 = \mathbb{RP}^2$. Its fiber over any point $x \in \mathbb{RP}^2$ is isomorphic to $\mathbb{R}^{2k}$ and consists of all linear systems vanishing on the corresponding line in $\mathbb{R}^3$. This fiber bundle is orientable if and only if k is even, so by Thom isomorphism we have $\overline{H}_i(\sigma_1, \mathbb{Q}) \simeq H_{i-2k}(\mathbb{RP}^2, \mathbb{Q})$ if k is even, and $\overline{H}_i(\sigma_1, \mathbb{Q}) \simeq H_{i-2k}(\mathbb{RP}^2, \mathbb{Q} \otimes \mathbb{O}r)$ if k is odd; here $\mathbb{O}r$ is the orientation bundle of $\mathbb{RP}^2$.

The space $\sigma_2 \setminus \sigma_1$ is fibered over $J_2 = \widehat{\mathbb{RP}^2}$. Its fiber over any point $l \in \widehat{\mathbb{RP}^2}$ is the product of the k-dimensional subspace $L(l) \subset \mathbb{R}^{3k}$ and

the open two-dimensional disc $\breve{\Psi}(l)$, which is the union of all half-open intervals connecting the point $l \in J_2$ with all points of J_1 corresponding to the points of the line l. The fiber bundle over J_2 with fibers $L(l)$ is orientable if and only if k is even; the fiber bundle with fibers $\breve{\Psi}(l)$ always is non-orientable. Therefore $\overline{H}_i(\sigma_2 \setminus \sigma_1, \mathbb{Q}) \simeq H_{i-k-2}(\mathbb{RP}^2, \mathbb{Q})$ if k is odd, and $\overline{H}_i(\sigma_2 \setminus \sigma_1, \mathbb{Q}) \simeq H_{i-k-2}(\mathbb{RP}^2, \mathbb{Q} \otimes \mathbb{O}r)$ if k is even. Finally, $\sigma_3 \setminus \sigma_2$ is in all cases the entire order complex Ψ of non-empty linear subspaces of $\mathbb{R}^3$, from which its order subcomplex $\partial\Psi$ of only proper subspaces is removed. Ψ is a cone over this subcomplex, which is known (see e.g. [6], [18]) to be homeomorphic to S^4. So, $\overline{H}_i(\sigma_3 \setminus \sigma_2, \mathbb{Q})$ is equal to $\mathbb{Q}$ if $i = 5$, and is trivial otherwise. Summarizing, we get that our spectral sequence has only the following non-trivial terms $E^1_{p,q}$ (all of which are equal to $\mathbb{Q}$). For even k this are terms with (p, q) equal to $(1, 2k-1), (2, k+2)$, and $(3, 2)$; for odd k the terms with (p, q) equal to $(1, 2k+1), (2, k)$, and $(3, 2)$ only. All further differentials of the spectral sequence are obviously trivial. We obtain the group $\overline{H}_*(\sigma, \mathbb{Q})$, and Proposition 1 follows by Alexander duality similar to (3) but with $6k$ replaced by $3k$. $\square$

2.3. **Singular sets for the resultant in the spaces of quadratic maps.** From now on, we come back to the space W^k of systems of k quadratic functions $\mathbb{R}^3 \to \mathbb{R}$, consider the resultant subvariety Σ in it, and construct the simplicial resolution σ of it. In this construction, we follow the A. Gorinov's (see [10], [15]) modification of the construction of resolutions of discriminant sets proposed in [20].

We fix the following list of families of compact subsets in $\mathbb{RP}^2$, basic for the construction of the resolution, cf. [20], [10].

$J_1 \sim \mathbb{RP}^2$ consists of particular points.

J_2 is the configuration space $B(\mathbb{RP}^2, 2)$ of unordered pairs of distinct points in $\mathbb{RP}^2$.

J_3 is the similar space $B(\mathbb{RP}^2, 3)$ of subsets of cardinality 3 in $\mathbb{RP}^2$.

J_4 is the space of quadruples of distinct points, all of which belong to one and the same line in $\mathbb{RP}^2$.

J_5 is the space $\widehat{\mathbb{RP}^2}$ of lines.

J_6 is the space $B^\times(\mathbb{RP}^2, 4)$ of quadruples of different points in $\mathbb{RP}^2$, which do not belong to one and the same line.

J_7 is the space of configurations of the form {a line, a point outside it in $\mathbb{RP}^2$}.

J_8 is the space $B(\widehat{\mathbb{RP}^2}, 2)$ of pairs of different lines in $\mathbb{RP}^2$.

J_9 is the space of non-singular non-empty conics in $\mathbb{RP}^2$.

J_{10} consists of unique point and corresponds to entire $\mathbb{RP}^2$.

Following [10], supply the union of these spaces with the Hausdorff metric on the space of compact subsets in $\mathbb{RP}^2$: the distance between two compact subsets $X, Y \subset \mathbb{RP}^2$ is

$$\max_{x\in X} \rho(x, Y) + \max_{y\in Y} \rho(y, X), \tag{4}$$

where ρ is the usual metric in the plane $\mathbb{RP}^2$ considered as a quotient space of the unit 2-sphere. Denote by $\mathfrak{J}$ the obtained metric space $J_1 \cup J_2 \cup \cdots \cup J_{10}$. It is easy to check that it is a closed subset of the space of all compact subsets of $\mathbb{RP}^2$ supplied with the metric (4). The latter space is compact, hence our space $\mathfrak{J}$ is compact too.

Again, consider this space $\mathfrak{J}$ as a partially ordered set by incidence of corresponding subsets. Unlike the situation considered in §2.2, some of its strata J_p are not compact and adjoin one another. By this reason the construction of the simplicial resolution is slightly more complicated. In particular, we use the following notion (see [3], [16]).

Definition 1. Given a compact finite-dimensional topological space X, let us embed it generically into a space $\mathbb{R}^\omega$ of a very large dimension, and define its *r-th self-join* X^{*r} as the union of all simplices of dimensions $1, \ldots, r-1$ in $\mathbb{R}^\omega$, all whose vertices belong to the image of X. (The genericity of an embedding means that intersections of different simplices are their common faces only).

It is easy to see that the spaces X^{*r} defined by different generic embeddings of X are canonically homeomorphic to one another.

Define the order complex Ψ of sets $J_1, \ldots, J_{10}$ as a subspace of the 10-th self-join $\mathfrak{J}^{*10}$: it consists of all possible simplices (of different dimensions) in $\mathfrak{J}^{*10}$, all whose vertices are incident to one another. For any $X \in \mathfrak{J}$, define the space $\Psi(X)$ as the union of simplices in Ψ, all whose vertices correspond to sets containing $\{X\}$. Also we define the link $\partial\Psi(X)$ as the union of simplices in $\Psi(X)$ not containing the point X, and the space $\breve{\Psi}(X)$ as the difference $\Psi(X) \setminus \partial\Psi(X)$. By the exact sequence of the pair we have

$$\overline{H}_i(\breve{\Psi}(X)) \equiv \tilde{H}_{i-1}(\partial\Psi(X)).$$

The space Ψ is naturally filtered: for any $p \in [1, 10]$ the space $\Psi_p \subset \Psi$ is defined as the union of spaces $\Psi(X)$ over all $X \in J_j$, $j \le p$. By the construction, $\Psi_p \setminus \Psi_{p-1}$ is the union of spaces $\breve{\Psi}(X)$ over all $X \in J_p$.

For any point $X \in J_p$ define the space $\sigma(X) \subset \Psi \times W^k$ as the direct product $\Psi(X) \times L(X)$, where $L(X)$ is the space of all systems vanishing on the set $\{X\}$. The space σ_p is defined as the union of spaces $\sigma(X)$

over all $X \in J_j$, $j \le p$; $\sigma \equiv \sigma_{10}$. So, we have filtrations

$$\Psi_1 \subset \cdots \subset \Psi_{10} \equiv \Psi, \qquad \sigma_1 \subset \cdots \subset \sigma_{10} \equiv \sigma. \tag{5}$$

The obvious projection $\Psi \times W^k \to W^k$ induces a map $\Pi : \sigma \to \Sigma$. For any point $F \in \Sigma$ (i.e. some resultant system of polynomials) its preimage $\Pi^{-1}(F)$ is the subcomplex $\Psi(X(F)) \times F$, where $X(F) \in \mathfrak{J}$ is the whole set of common zeros of all polynomials from F. (Any such set belongs to some J_p).

Lemma 1. *1. The map $\Pi : \sigma \to \Sigma$ induced by the obvious projection $\Psi \times W^k \to W^k$ is proper, and its continuation to a map of one-point compactifications of these spaces σ and Σ is a homotopy equivalence. In particular, it induces an isomorphism of Borel–Moore homology groups, $\overline{H}_*(\sigma) \simeq \overline{H}_*(\Sigma)$.*

2. For any $p = 1, \ldots, 10$, the projection $\sigma \to \Psi$ maps σ_p to Ψ_p. Its restriction to $\sigma_p \setminus \sigma_{p-1}$ is a vector bundle over $\Psi_p \setminus \Psi_{p-1}$, whose dimension is equal to that of the space $L(X)$ for any $X \in J_p$.

3. For any p, $\Psi_p \setminus \Psi_{p-1}$ is locally trivially fibered over J_p, with the fiber over $X \in J_p$ equal to $\breve{\Psi}(X)$.

For the proof of item 1, see [10], Theorem 2.8. Indeed, our collection of families $J_1, \ldots, J_{10}$ obviously satisfies all conditions listed in [10], p. 399 (see also [15], pp. 13–14). Items 2 and 3 follow immediately from the construction. $\square$

So, we can (and will) calculate the group $\overline{H}_*(\sigma)$ instead of $\overline{H}_*(\Sigma)$.

Consider the spectral sequence $E^r_{p,q}$, calculating the group $\overline{H}_*(\sigma, \mathbb{Q})$ and generated by the filtration (5), with unique reservation: we do not distinguish σ_4 from the rest of σ_5, so that the revised filtration looks as follows: $\sigma_1 \subset \sigma_2 \subset \sigma_3 \subset \sigma_5 \subset \ldots$. The term $E^1_{p,q}$ is thus canonically isomorphic to the group $\overline{H}_{p+q}(\sigma_p \setminus \sigma_{p-1}, \mathbb{Q})$ if $p = 1, 2$ or 3, to $\overline{H}_{4+q}(\sigma_5 \setminus \sigma_3, \mathbb{Q})$ if $p = 4$, and $\overline{H}_{p+q}(\sigma_{p+1} \setminus \sigma_p, \mathbb{Q})$ for $p = 5, \ldots, 9$.

Theorem 2. *For any even (respectively, odd) k, all non-trivial groups $E^1_{p,q}$ of the initial term of our spectral sequence are as shown in Fig. 1 left (respectively, right). In the case of odd k, the differential $d^1 : E^1_{8,k+5} \to E^1_{7,k+5}$ is an isomorphism.*

Theorem 1 for $k \ge 3$ follows immediately from this one, because no other differentials $d^r : E^r_{p,q} \to E^r_{p-r,q+r-1}$ of the spectral sequence can connect non-trivial cells of these tables.

For $k = 2$ the left-hand table of Fig. 1 becomes as is shown in Fig. 2. Since the entire space W^2 is 12-dimensional, all groups $E_{p,q}$ with



FIG. 1. E^1 for k even (k=6) and k odd (k=5)



FIG. 2. E^1 for $k = 2$

$p + q \geq 12$ should be killed by some differentials. This can happen in unique way: the differentials $d^4 : E^4_{8,7} \to E^4_{4,10}$ and $d^3 : E^3_{9,5} \to E^3_{6,7}$ should be isomorphisms. Thus, the only surviving term is $E_{1,9} \simeq \mathbb{Q}$.

This gives us $\overline{H}_{10}(\Sigma, \mathbb{Q}) \simeq \mathbb{Q}$ and proves (via the Alexander duality) the assertion of Theorem 1 for the remaining case $k = 2$.

The proof of Theorem 2 takes the rest of this article.

Remark 1. In both spectral sequences shown in Fig. 1, the column $p = 9$ consists of the Borel–Moore homology groups of $\sigma_{10} \setminus \sigma_9 \equiv \Psi_{10} \setminus \Psi_9 \equiv \breve{\Psi}([\mathbb{RP}^2])$. Namely, $E^1_{9,q} \simeq \overline{H}_{9+q}(\breve{\Psi}([\mathbb{RP}^2]), \mathbb{Q}) \equiv \tilde{H}_{8+q}(\partial\Psi([\mathbb{RP}^2]), \mathbb{Q})$. The latter space $\partial\Psi([\mathbb{RP}^2])$ is the simplicial resolution of the topological order complex of the partially ordered set of all proper subsets in $\mathbb{RP}^2$, listed in §2.3. Thus we prove in particular that this order complex has rational homology group of S^{13}. This order complex is very similar (but does not coincide) with the order complex of all proper subspaces in W consisting of quadrics vanishing on some subset in $\mathbb{RP}^2$. The latter complex is studied in more detail in [22]. In particular, it is proved there that this complex also has rational homology of S^{13}.

Notice that the similar order complex of sets defined by systems of linear functions in $\mathbb{R}^3$ is homeomorphic to S^4; we have used this fact in the proof of Proposition 1.

3. Preliminary facts on homology of configuration spaces

Given a topological space M, denote by $B(M, j)$ the *configuration space*, whose points are the subsets of cardinality j in M.

Lemma 2. *For any j, there is a locally trivial fibre bundle $B(S^1, j) \to S^1$, whose fiber is homeomorphic to $\mathbb{R}^{j-1}$. This fibre bundle is orientable (and hence trivial) if j is odd and is non-orientable if j is even.*

Indeed, consider S^1 as the unit circle in $\mathbb{C}^1$, then the projection map of this fibre bundle can be realised as taking the product of j complex numbers on this circle. The fibre of this bundle can be identified in the terms of the universal covering $\mathbb{R}^j \to T^j$ with any connected component of some hyperplane $\{x_1 + \cdots + x_j = \text{const}\} \subset \mathbb{R}^j$, from which all affine planes given by $x_m = x_n + 2\pi k$, $m \neq n$, $k \in \mathbb{Z}$, are removed. Any such component is convex and hence diffeomorphic to $\mathbb{R}^{j-1}$. The assertion on the orientability can be checked immediately. □

Remark 2. The image of the standard section of this fiber bundle consists of all j-ples of points separating S^1 into equal arcs.

For any manifold M, we denote by $\mathbb{O}r$ the orientation local system of groups on M: it is locally isomorphic to the constant $\mathbb{Z}$-system, but the elements of $\pi_1(M)$ violating the orientation of M act on it as the multiplication by -1. For any topological space M denote by $\pm\mathbb{Z}$ the

local system on $B(M,j)$, which is locally isomorphic to the constant $\mathbb{Z}$-bundle, but the elements of $\pi_1(B(M,j))$ defining odd permutations of j points act on it as multiplication by -1. Also, set $\pm\mathbb{Q} \equiv \pm\mathbb{Z} \otimes \mathbb{Q}$.

Lemma 3. *For any even-dimensional manifold M, a closed loop in the manifold $B(M,j)$ violates the orientation of this manifold if and only if the union of traces of all j points of our configuration during their corresponding movement defines a disorienting cycle in $H_1(M,\mathbb{Z}_2)$ (i.e. a cycle taking non-zero value on the first Stiefel–Whitney class of M).*

Proof is trivial, see e.g. [17]. □

Consider the $j!$-fold covering $\nu : I(\mathbb{RP}^2,j) \to B(\mathbb{RP}^2,j)$, where $I(\mathbb{RP}^2,j)$ is the space of *ordered* sets of j distinct points in $\mathbb{RP}^2$.

Lemma 4. *For any natural $j \geq 2$ the Borel–Moore homology group $\overline{H}_*(I(\mathbb{RP}^2,j),\mathbb{Q})$ is trivial in all dimensions.*

Proof. $I(\mathbb{RP}^2,j) = (\mathbb{RP}^2)^j \setminus \Delta$, where $\Delta \subset (\mathbb{RP}^2)^j$ is the set of all sequences of j points in $\mathbb{RP}^2$, at least two of which coincide. By the exact sequence of the pair $((\mathbb{RP}^2)^j,\Delta)$ it is enough to prove that $\tilde{H}_*(\Delta,\mathbb{Q}) = 0$ in all dimensions. Δ is the union of $\binom{j}{2}$ spaces $\Delta_{r,s} \simeq (\mathbb{RP}^2)^{j-1}$, $1 \leq r < s \leq j$, defined by the coincidence of the r-th and s-th members of the sequences of j points in $\mathbb{RP}^2$. Consider the Mayer-Vietoris spectral sequence of this union. All (maybe, multiple) intersections of spaces $\Delta_{r,s}$ are homeomorphic to $(\mathbb{RP}^2)^m$, $m \in [1, j-1]$, in particular have the rational homology of a point. Therefore this spectral sequence reduces to the one calculating the homology of the order complex of all such multiple intersections. This order complex has a maximal element (corresponding to the non-empty intersection of all these spaces), hence is contractible. □

Corollary 1. *For any $j \geq 2$, the groups $\overline{H}_*(B(\mathbb{RP}^2,j),\mathbb{Q})$ and $\overline{H}_*(B(\mathbb{RP}^2,j),\pm\mathbb{Q})$ are trivial in all dimensions.*

Indeed, $\overline{H}_*(I(\mathbb{RP}^2,j),\mathbb{Q}) \simeq \overline{H}_*(B(\mathbb{RP}^2,j),\nu_!(\mathbb{Q}))$, where $\nu_!(\mathbb{Q})$ is the direct image of the constant $\mathbb{Q}$-sheaf on $I(\mathbb{RP}^2,j)$. $\nu_!(\mathbb{Q})$ is a $j!$-dimensional local system on $B(\mathbb{RP}^2,j)$, containing as direct summands both local systems $\mathbb{Q}$ and $\pm\mathbb{Q}$. So, both groups mentioned in Corollary 1 are direct summands of the group $\overline{H}_*(I(\mathbb{RP}^2,j),\mathbb{Q})$, which is trivial by Lemma 4. □

Recall the notation $B^\times(\mathbb{RP}^2,4)$ for the subset in $B(\mathbb{RP}^2,4)$ consisting of collections not contained entirely in any line in $\mathbb{RP}^2$.

Lemma 5. *The Borel–Moore homology groups $\overline{H}_*(B^\times(\mathbb{RP}^2,4),\mathbb{Q})$ and $\overline{H}_*(B^\times(\mathbb{RP}^2,4),\pm\mathbb{Q})$ are trivial in all dimensions.*

Proof. Denote by T the difference $B(\mathbb{RP}^2,4)\setminus B^\times(\mathbb{RP}^2,4)$. By Corollary 1 and exact sequence of the pair $(B(\mathbb{RP}^2,4),T)$, it is enough to prove that the groups $\overline{H}_*(T,\mathbb{Q})$ and $\overline{H}_*(T,\pm\mathbb{Q})$ are trivial. By Poincaré duality, this is equivalent to the triviality of groups $H_*(T,\mathbb{Q}\otimes\mathbb{O}r)$ and $H_*(T,\pm\mathbb{Q}\otimes\mathbb{O}r)$.

T is fibered over the space $\widehat{\mathbb{RP}^2}$ of all lines in $\mathbb{RP}^2$. Its fiber over the point, corresponding to the line l, is the configuration space $B(l,4)$. By Lemma 2 this fiber is homotopy equivalent to S^1. The retractions of these fibers to the circles described in Remark 2 can be performed uniformly, therefore T contains a 3-dimensional deformation retract $\tilde{T}$ also fibered over $\mathbb{RP}^2$; its fiber over the projective line l consists of all quadruples of points separating this line (considered as a circle in the unit sphere factored by the central symmetry) into arcs of length $\pi/4$. So, it remains to prove the triviality of groups

$$H_*(\tilde{T},\mathbb{Q}\otimes\widetilde{\mathbb{O}r}) \qquad \text{and} \qquad H_*(\tilde{T},\pm\mathbb{Q}\otimes\widetilde{\mathbb{O}r}), \tag{6}$$

where $\widetilde{\mathbb{O}r}$ is the restriction to $\tilde{T}$ of the orientation sheaf of T.

The group $SO(3)$ (and hence also it double cover $SU(2)$) acts transitively on $\tilde{T}$; this action defines a principal 16-fold covering $SU(2)\to\tilde{T}$. The 16-dimensional direct image of the trivial $\mathbb{Q}$-bundle under this covering contains both sheaves $\mathbb{Q}\otimes\widetilde{\mathbb{O}r})$ and $\pm\mathbb{Q}\otimes\widetilde{\mathbb{O}r}$ as direct summands, hence both groups (6) are at most 1-dimensional in dimensions 0 and 3 and are trivial in all other dimensions. However, both these 1-dimensional sheaves are not equivalent to the constant one, therefore the groups (6) are trivial in dimension 0; by the Euler characteristic considerations the same is true for these groups in dimension 3. □

Lemma 6. *The group $\overline{H}_i(B(\mathbb{RP}^2,2),\mathbb{O}r\otimes\mathbb{Q})$ is equal to $\mathbb{Q}$ for $i=4$ and $i=1$, and is trivial in all other dimensions.*

Proof. This group is Poincaré dual to $H_{4-i}(B(\mathbb{RP}^2,2),\mathbb{Q})$; let us calculate the latter group. Associating with any two-point configuration the line spanned by these points, we obtain the fibration of $B(\mathbb{RP}^2,2)$ over $\mathbb{RP}^2$. Its fiber is homotopy equivalent to S^1. The generator of $\pi_1(\mathbb{RP}^2)$ acts non-trivially on the group H_1 of the fiber, thus the term E^2 of the homological spectral sequence of this fibration has only two non-trivial cells, $E^2_{0,0}\simeq\mathbb{Q}$ and $E^2_{2,1}\simeq H_2(\mathbb{RP}^2,\mathbb{O}r\otimes\mathbb{Q})\simeq\mathbb{Q}$. □

Proposition 2 (Caratheodory theorem, see e.g. [18], [8])**.** *For any $r\geq 1$, the r-th self-join $(S^1)^{*r}$ of S^1 is homeomorphic to S^{2r-1}.* □

Lemma 7. *If r is even, then the sphere $(S^1)^{*r}\sim S^{2r-1}$ has a canonical orientation, and hence a canonical generator of its $(2r-1)$-dimensional*

homology group. If r is odd, then the homeomorphism of $(S^1)^{*r}$ to itself, induced by any orientation-reversing automorphism of S^1, violates the orientations of $(S^1)^{*r}$.

Proof. The group $H_{2r-1}((S^1)^{*r})$ is generated by the Borel–Moore fundamental class of an open dense subset in $(S^1)^{*r}$, which is fibered over the configuration space $B(S^1, r)$. Its fiber over an r-configuration is the open $(r-1)$-dimensional simplex, whose vertices correspond to the points of this configuration. The orientation of this space consists of the orientations of the base and the fibers. Given an r-configuration in S^1, let us number arbitrarily its points, and define the j-th tangent vector of the base, $j = 1, \dots, r$, as a shift of the j-th point along the chosen orientation of S^1, keeping all remaining $r-1$ points unmoved. The fiber over this configuration will be oriented by the same order of these r points. Renumbering these points, we simultaneously change or preserve the orientations of the base and the fibre, thus preserving the orientation of the total space. On the other hand, changing the chosen orientation of S^1 we reverse r basic vectors of the orienting frame. □

4. Calculation of the spectral sequence

By item 2 of Lemma 1 and the Thom isomorphism, we have

$$\overline{H}_m(\sigma_p \setminus \sigma_{p-1}, \mathbb{Q}) \simeq \overline{H}_{m-d(p)\cdot k}(\Psi_p \setminus \Psi_{p-1}, \mathbb{Q} \otimes \mathrm{O}) \tag{7}$$

for any m and p; here O is the orientation sheaf of the vector bundle, whose fiber over any point of $\breve{\Psi}(X)$, $X \in J_p$, is the space $L(X) \subset W^k$ of all quadratic systems vanishing on the set $\{X\}$, and $d(p)$ is the dimension of these spaces $L(X), X \in J_p$ (i.e. $d(1) = 5$, $d(2) = 4$, $d(3) = d(4) = d(5) = 3$, $d(6) = d(7) = 2$, $d(8) = d(9) = 1$, and $d(10) = 0$). By the construction, this vector bundle is the direct sum of k copies of a bundle, whose fibers are corresponding subspaces of W. Therefore this orientation sheaf is trivial if k is even, and coincides with the orientation sheaf of the latter vector bundle if k is odd.

So, we are going to calculate the right-hand groups in (7).

4.1. **First column.** The first term σ_1 of our filtration of σ is the space of a $5k$-dimensional vector bundle over $\mathbb{RP}^2$. This vector bundle is naturally oriented for any k. Indeed, any of its k summands is a 5-dimensional vector subbundle of the trivial bundle $\mathbb{RP}^2 \times W \to \mathbb{RP}^2$; its fiber over a point $x \in \mathbb{RP}^2$ is the space of all polynomials $f \in W^k$ vanishing on the line $\{x\} \subset \mathbb{R}^3$. The canonical coorientation of such a fiber in W is defined by the increase of the polynomials f on the

corresponding line $\{x\}$. By (7) we have

$$E^1_{1,q} \equiv \overline{H}_{1+q-5k}(\mathbb{RP}^2, \mathbb{Q}). \tag{8}$$

So, the group $E^1_{1,q}$ is equal to $\mathbb{Q}$ if $q = 5k-1$ and is trivial otherwise.

4.2. **Second column.**

Lemma 8. *The group $\overline{H}_*(\sigma_2 \setminus \sigma_1, \mathbb{Q})$ is trivial in all dimensions.*

Proof. $\sigma_2 \setminus \sigma_1$ is the space of a fiber bundle over the configuration space $B(\mathbb{RP}^2, 2)$. Namely, it is the fibered product of

— the $(4k)$-dimensional vector bundle, whose fiber over the point $\{x, y\} \in B(\mathbb{RP}^2, 2)$ consists of all systems $F \in W^k$ vanishing on both lines in $\mathbb{R}^3$ corresponding to the points x and y, and

— the bundle of open intervals, whose endpoints are associated with the points x and y. Any such interval is constituted by two half-intervals $(\{x\}, \{x \cup y\}]$ and $(\{y\}, \{x \cup y\}]$ in $\breve{\Psi}_2$, where $\{x\}$ and $\{y\}$ are points of J_1 corresponding to the points x and y, and $\{x \cup y\} \in J_2$. The endpoints $\{x\}$ and $\{y\}$ are excluded from the segment as they belong to the first term Ψ_1 of our filtration.

Any loop in $B(\mathbb{RP}^2, 2)$ not permuting two points of the configuration does not violate the orientations of these two bundles. A loop permuting these points changes the orientations of all k factors of the first bundle, as well as the orientation of the second. Thus the m-dimensional Borel–Moore homology group of the total space of this fiber bundle is equal to $\overline{H}_{m-4k-1}(B(\mathbb{RP}^2, 2), \mathbb{Q})$ if k is odd, and to $\overline{H}_{m-4k-1}(B(\mathbb{RP}^2, 2), \pm\mathbb{Q})$ if k is even. By Corollary 1 of Lemma 4 all these groups are trivial. □

Therefore the column $E^1_{2,*}$ of the main spectral sequence consists of zeroes only.

4.3. **Third column.** The space $\sigma_3 \setminus \sigma_2$ is fibered over the configuration space $J_3 = B(\mathbb{RP}^2, 3)$. Its fiber over a 3-point configuration $X \in J_3$ is equal to the product of the $3k$-dimensional vector space $L(X)$ and the *open* 2-dimensional simplex (whose vertices are associated with the points of the configuration, and the entire simplex is the order complex of non-empty subsets of this configuration). Any element of $\pi_1(B(\mathbb{RP}^2, 3))$, defining an odd permutation of three points, changes the orientations of the bundle of triangles and of all k summands of the $3k$-dimensional vector bundle. Therefore we have the following lemma.

Lemma 9. *For even k, $\overline{H}_m(\sigma_3 \setminus \sigma_2, \mathbb{Q}) = \overline{H}_{m-3k-2}(B(\mathbb{RP}^2, 3), \pm\mathbb{Q})$. For odd k, $\overline{H}_m(\sigma_3 \setminus \sigma_2, \mathbb{Q}) = \overline{H}_{m-3k-2}(B(\mathbb{RP}^2, 3), \mathbb{Q})$.*

By Corollary 1 of Lemma 4 all these groups are trivial in all dimensions, so the column $p = 3$ of the main spectral sequence is trivial.

4.4. **Fourth column.** The space $\sigma_5 \setminus \sigma_3$ is fibered over the space $\widehat{\mathbb{RP}^2}$ of all lines in $\mathbb{RP}^2$. Its fiber over the line l is equal to the product of the $3k$-dimensional subspace $L(l) \subset W^k$ (consisting of all polynomial systems vanishing on l) and the space $\Psi(l) \setminus \Psi_3$.

The space $\Psi(l)$ is a compact cone with the vertex $\{l\} \in J_5$, in particular is contractible. Its subset $\Psi(l) \cap \Psi_3$ is the order complex of all subsets of cardinality ≤ 3 in the line $l \sim S^1$. It is the union of order subcomplexes $\Psi(\{a, b, c\})$, where a, b, c are arbitrary distinct points of l. Any such subcomplex can be identified with the triangle with vertices $\{a\}, \{b\}$ and $\{c\}$; the entire space $\Psi(l) \cap \Psi_3$ becomes thus identified with the union of all such simplices supplied with the natural topology. By Proposition 2 this space is homeomorphic to S^5. The exact sequence of the pair $(\Psi(l), \Psi(l) \cap \Psi_3)$ proves now that the Borel–Moore homology of the fiber $\Psi(l) \setminus \Psi_3$ is equal to $\mathbb{Q}$ in dimension 6 and is trivial in all other dimensions. By Lemma 7 the monodromy over $\widehat{\mathbb{RP}^2}$ reverses the orientation of this factor of the fiber, and also of this homology group.

On the other hand, the bundle with fiber $L(l)$ splits into the direct sum of k 3-dimensional bundles, any of which changes its orientation over the generator of $\pi_1(\widehat{\mathbb{RP}^2})$. So, we obtain the following fact.

Lemma 10. *For even k, $\overline{H}_i(\sigma_5 \setminus \sigma_3, \mathbb{Q}) \simeq \overline{H}_{i-3k-6}(\widehat{\mathbb{RP}^2}, \mathbb{Q} \otimes \mathbb{O}r)$. In particular, it is equal to $\mathbb{Q}$ if $i = 3k + 8$ and is trivial in all other dimensions.*

For odd k, $\overline{H}_i(\sigma_5 \setminus \sigma_3, \mathbb{Q}) \simeq \overline{H}_{i-3k-6}(\widehat{\mathbb{RP}^2}, \mathbb{Q})$. In particular, it is equal to $\mathbb{Q}$ if $i = 3k + 6$ and is trivial in all other dimensions. □

This gives us the columns $p = 4$ of both parts of Fig. 1. □

4.5. **Fifth column.** The space $\sigma_6 \setminus \sigma_5$ is fibered over the restricted configuration space $B^\times(\mathbb{RP}^2, 4)$. Its fiber over $X \in B^\times(\mathbb{RP}^2, 4)$ is the product of the $2k$-dimensional vector space $L(X)$ and the open 3-dimensional simplex, whose vertices correspond to the points of the collection $\{X\}$. In exactly the same way as in §§4.2, 4.3, we obtain the following fact.

Lemma 11. *The group $\overline{H}_m(\sigma_6 \setminus \sigma_5, \mathbb{Q})$ is equal to $\overline{H}_{m-2k-3}(B^\times(\mathbb{RP}^2, 4), \mathbb{Q})$ if k is odd, and to $\overline{H}_{m-2k-3}(B^\times(\mathbb{RP}^2, 4), \pm\mathbb{Q})$ if k is even.*

By Lemma 5 all these groups are trivial for any m, and the column $p = 5$ of our spectral sequence is zero.

4.6. **Sixth column.**

Lemma 12. *For odd k, $\overline{H}_i(\sigma_7 \setminus \sigma_6, \mathbb{Q}) \simeq \overline{H}_{i-2k-9}(\mathbb{RP}^2, \mathbb{Q} \otimes \mathbb{O}r)$, in particular this group is equal to $\mathbb{Q}$ if $i = 2k+11$ and is trivial otherwise. For even k, $\overline{H}_i(\sigma_7 \setminus \sigma_6, \mathbb{Q}) \simeq \overline{H}_{i-2k-9}(\mathbb{RP}^2, \mathbb{Q})$, in particular is equal to $\mathbb{Q}$ if $i = 2k + 9$ and is trivial otherwise.*

Proof. $\sigma_7 \setminus \sigma_6$ is the space of a fiber bundle over the space J_7 of configurations

$$\{\text{a line } \{l\} \in \widehat{\mathbb{RP}^2}; \text{a point } u \in \mathbb{RP}^2 \setminus l\}. \tag{9}$$

Its fiber over such a configuration $X = \{l, u\} \in J_7$ is the product

$$\breve{\Psi}(X) \times L(X), \tag{10}$$

where $L(X) \subset W^k$ is the $2k$-dimensional space of all systems $F \in W^k$ vanishing on l and on u. The space of configurations (9) is obviously diffeomorphic to $T_*\mathbb{RP}^2$, in particular is orientable.

Lemma 13. *1. The non-trivial element of the fundamental group $\pi_1(T_*\mathbb{RP}^2) \simeq \pi_1(\mathbb{RP}^2)$ of this configuration space violates the orientation of any of k two-dimensional summands of the second factor $L(X)$ of the fiber (10).*

2. The link $\partial\Psi(X)$ of the first factor of the fiber (10) is homotopy equivalent to S^6, in particular the group $\overline{H}_i(\breve{\Psi}(X), \mathbb{Q})$ is isomorphic to $\mathbb{Q}$ if $i = 7$ and is trivial for all other i.

3. The monodromy over the non-trivial element of $\pi_1(T_*\mathbb{RP}^2)$ acts as multiplication by -1 on the homology bundle associated with the latter factors.

Proof. 1. For the loop in our configuration space, generating its fundamental group, we choose the family of pairs $\{l_\varphi, u_\varphi\}$, $\varphi \in [0, \pi]$, where the line l_φ is defined by the equation $x\cos\varphi + y\sin\varphi = 0$, and the point u_φ by homogeneous coordinates $(\cos\varphi : \sin\varphi : 0)$. For any configuration $X = \{l, u\}$, the fiber of any of our k summands over X consists of all quadratic forms vanishing on l and u. For two independent sections of the two-dimensional bundle of such forms over the set of pairs $\{l_\varphi, u_\varphi\}$, where φ runs over the half-interval $\varphi \in [0, \pi)$, we choose the families

$$(x\cos\varphi + y\sin\varphi)(x\sin\varphi - y\cos\varphi) \quad \text{and} \quad (x\cos\varphi + y\sin\varphi)z.$$

The first of these sections extends to a continuous section over the entire closed loop, while the limit value of the second one at $\varphi = \pi$ is opposite to the value at $\varphi = 0$. So, the monodromy over this loop acts on the fiber as the operator $\begin{vmatrix} 1 & 0 \\ 0 & -1 \end{vmatrix}$.

2. The link $\partial\Psi(X)$, $X = \{l, u\}$, consists of two subspaces: $A =$ the order complex $\Psi(l)$, and $B =$ the union of all order complexes $\Psi(\{a, b, c, u\})$, where a, b and c are arbitrary distinct points in the line l. The intersection $A \cap B$ is the space $\Psi(l) \cap \Psi_3$, i.e. the union of similar order subcomplexes $\Psi(\{a, b, c\})$. Since §4.4 we know that this intersection $A \cap B$ is homeomorphic to S^5. The set A is compact and contractible, hence the pair $(A, A \cap B)$ is homotopy equivalent to the pair (cone over $A \cap B$, its base $A \cap B$). As for the space B, any of order complexes $\Psi(\{a, b, c, u\})$ constituting it is canonically PL-homeomorphic to the tetrahedron with vertices a, b, c and u, and hence to the cone over $\Psi(\{a, b, c\})$ with vertex $\{u\}$. These homeomorphisms over all triples $\{a, b, c\}$ define a homeomorphism between the union B of all these tetrahedra and another cone over $\Psi(l) \cap \Psi_3$. Therefore our space $A \cup B$ is homotopy equivalent to the suspension over $A \cap B \sim S^5$.

3. By Lemma 7, the generator of $\pi_1(\mathbb{RP}^2)$ acts on the group $H_5(\Psi(l) \cap \Psi_3, \mathbb{Q})$ as multiplication by -1. Two parts A and B of $\partial\Psi(\{l, u\}) \sim \Sigma(\Psi(l) \cap \Psi_3)$ are of different origin and cannot be permuted by the monodromy, hence the group $H_6(\partial\Psi(\{l, u\}), \mathbb{Q})$ also is reversed by this generator. □

Lemma 13 is completely proved, and Lemma 12 follows immediately from it by a) the spectral sequence of the fiber bundle (10) over the space $J_6 \sim T_*\mathbb{RP}^2$ of configurations (9), and b) the Thom isomorphism for the non-orientable fibration of this space over $\mathbb{RP}^2$. □

This gives us columns $p = 6$ of both spectral sequences of Fig. 1.

4.7. **Seventh column.**

Proposition 3. *For any even k, the group $\overline{H}_*(\sigma_8 \setminus \sigma_7, \mathbb{Q})$ is trivial in all dimensions. If k is odd, then $\overline{H}_i(\sigma_8 \setminus \sigma_7, \mathbb{Q})$ is isomorphic to $\mathbb{Q}$ for $i = k + 12$ and $i = k + 9$, and is trivial for all other values of i.*

Proof. The space $\sigma_8 \setminus \sigma_7$ is fibered over the space $J_8 = B(\widehat{\mathbb{RP}^2}, 2)$ of 2-point configurations in the dual projective space; its fiber over any point X of J_8 is homeomorphic to $\breve{\Psi}(X) \times \mathbb{R}^k$.

Lemma 14. *For any point $X \in B(\widehat{\mathbb{RP}^2}, 2)$, the link $\partial\Psi(X)$ of the order complex $\Psi(X)$ is homology equivalent to the sphere S^7.*

Proof. Denote by l and l' two lines constituting the configuration X, and by O their common point. Let us filter the link $\partial\Psi(X)$:

$$(\Psi(l) \cup \Psi(l')) \subset (\partial\Psi(X) \cap \Psi_6) \subset \partial\Psi(X), \tag{11}$$

and study this filtration.

The order complexes $\Psi(l)$ and $\Psi(l')$ are contractible by definition, and their intersection set is a single point in Ψ_1, corresponding to the point $l \cap l'$. Therefore the union $\Psi(l) \cup \Psi(l')$ is contractible too.

The space $(\Psi(X) \cap \Psi_6) \setminus (\Psi(l) \cup \Psi(l'))$ consists of two similar disjoint pieces, swept out by all order complexes $\Psi(\{l, a'\}) \setminus \Psi(l)$, (respectively, $\Psi(\{l', a\}) \setminus \Psi(l')$,) where a' runs over the affine line $l' \setminus O$ and a runs over $l \setminus O$. Obviously, the first piece is homeomorphic to the direct product $(l' \setminus O) \times (\Psi(\{l, a'\}) \setminus \Psi(l))$ for an arbitrary $a' \in l' \setminus O$. By the proof of Lemma 13(2), the pair $(\Psi(\{l, a'\}), \Psi(l))$ is homotopy equivalent to the pair (D^7, D^6), where D^7 is the 7-dimensional ball, and D^6 is a closed hemisphere of its boundary $\partial D^7 \sim S^6$. In particular, the group $H_m(\Psi(\{l, a'\}), \Psi(l)) \equiv \overline{H}_m(\Psi(\{l, a'\}) \setminus \Psi(l))$ is trivial for any m. Hence by the Künneth formula all Borel-Moore homology groups of the first piece of $(\Psi(X) \cap \Psi_6) \setminus (\Psi(l) \cup \Psi(l'))$ also are trivial. The second piece is homeomorphic to it. So, the entire second term of the filtration (11) also has the homology of a point.

Finally, the space $\partial\Psi(X) \setminus \Psi_6$ is the union of all spaces $\breve{\Psi}(\{a, b, c, d\})$ where $a \neq b \in l \setminus O$, and $c \neq d \in l' \setminus O$. These spaces are homeomorphic to an open tetrahedron, and the base of this family, $B(\mathbb{R}^1, 2) \times B(\mathbb{R}^1, 2)$ is homeomorphic to the 4-dimensional open ball. In particular, the Borel–Moore homology group of this space is equal to $\mathbb{Q}$ in dimension 7 and trivial in all other dimensions. $\square$

Now, let us study the action of $\pi_1(B(\widehat{\mathbb{RP}^2}, 2))$ on the Borel–Moore homology group of fibers $\breve{\Psi}(X) \times L(X)$ of our fibration of $\sigma_7 \setminus \sigma_6$ over $B(\widehat{\mathbb{RP}^2}, 2)$. By the previous lemma this group is equal to $\mathbb{Q}$ in dimension $8 + k$ and is trivial in all other dimensions.

Let us choose for the basepoint in $B(\widehat{\mathbb{RP}^2}, 2)$ the pair of lines, one of which is obtained from meridians with longitudes 0 and 180, and the other with longitudes 90 and 270. The group $\pi_1(B(\widehat{\mathbb{RP}^2}, 2))$ is generated by two elements, θ and ϖ. Namely, θ is defined by the 90° rotation of the sphere around the main axis containing the poles; ϖ rotates the first line by 180° around the axis connecting the equator point with coordinates $(0, 0)$ and $(0, 180)$, and keeps the second line unmoved.

Lemma 15. *Both loops $\theta, \varpi \in \pi_1(B(\widehat{\mathbb{RP}^2}, 2))$ change the orientation of $B(\widehat{\mathbb{RP}^2}, 2)$ and of any of k factors of the fiber bundle with fibers $L(X)$. The one-dimensional homology groups $\overline{H}_8(\breve{\Psi}(X), \mathbb{Q})$ and*

$H_7(\partial\Psi(X),\mathbb{Q})$ have a canonical orientation for all X, depending continuously on X; in particular the group $\pi_1(B(\widehat{\mathbb{RP}^2},2))$ acts trivially on them.

Proof. The statement on the orientation of the base follows immediately from Lemma 3. Any of k summands of the fiber $L(X)$ is the space of quadratic functions vanishing on our pair of lines. We can choose one of two domains into which these lines divide $\mathbb{RP}^2$, and orient any summand by the increase of our quadratic functions inside this domain. The continuation along any of loops θ or ϖ permutes these two domains, hence violates this orientation.

Now we define an invariant orientation of the homology bundle whose fiber over $X \in B(\widehat{\mathbb{RP}^2},2)$ is the group $H_7(\partial\Psi(X),\mathbb{Q})$. This group is generated by the fundamental class of the 7-dimensional cell described in the last paragraph of the proof of Lemma 14. To orient this cell, we need to choose the orientations of its base $B(l\setminus O,2)\times B(l'\setminus O,2)$ and, for any point of this base (i.e. a configuration of four points), of the fiber over it, i.e. the 3-dimensional simplex, whose four vertices correspond to the points of this configuration. To do it, we fix an arbitrary numbering of these four points, and define the jth basic tangent vector to the base, $j=1,\dots,4$, as a shift of the jth point in the direction from the other point of the configuration in the same line and towards the intersection point O of these lines. The orientation of the fiber (i.e. a tetrahedron) is defined by the same numbering of its vertices. A different numbering of four points simultaneously changes or preserves the orientations of the base and the fiber, and hence preserves the orientation of the total space.

The orientation thus defined is obviously invariant and defines a trivialization of the homology bundle over $B(\widehat{\mathbb{RP}^2},2)$ with the fiber $H_7(\partial\Psi(X),\mathbb{Q})$ or, which is the same by the boundary isomorphism, of the group $\overline{H}_8(\breve{\Psi}(X),\mathbb{Q})$. □

Corollary 2. *If k is even (respectively, odd), then $\overline{H}_i(\sigma_7\setminus\sigma_6,\mathbb{Q})$ is equal to $\overline{H}_{i-k-8}(B(\widehat{\mathbb{RP}^2},2),\mathbb{Q})$ (respectively, $\overline{H}_{i-k-8}(B(\widehat{\mathbb{RP}^2},2),\mathbb{O}r\otimes\mathbb{Q})$) for any i.*

This fact follows from Lemma 15 by the Thom isomorphism. □

Proposition 3 follows from it by Corollary 1 and Lemma 6, and justifies the columns $p=7$ in both parts of Fig. 1. □

4.8. **Eighth column.**

Lemma 16. *The space J_9 of non-empty non-singular conics in $\mathbb{R}^3$ is diffeomorphic to the space of a 3-dimensional vector bundle over $\mathbb{RP}^2$, namely of the symmetric square of the cotangent bundle $S^2T^*(\mathbb{RP}^2)$. In particular, it is an orientable 5-dimensional manifold homotopy equivalent to $\mathbb{RP}^2$.*

Proof. Any such conic can be specified uniquely (up to the positive scalings) by a quadratic form of signature $(1, 2)$. Let us fix an arbitrary Euclidean structure in $\mathbb{R}^3$. Then the line of positive eigenvectors of these quadratic forms indicates a point of $\mathbb{RP}^2$, and the plane of negative ones can be naturally identified with the tangent plane of $\mathbb{RP}^2$ at this point. Let us always choose our quadratic form in such a way that it takes value 1 on its positive eigenvectors of length 1. Then we obtain an isomorphism between the space of all conics of this signature with this positive eigenvector on one hand, and the space of negative quadratic forms on the tangent space of $\mathbb{RP}^2$ at this point u on the other. The latter space is a convex open subset in the space $S^2T^*_u(\mathbb{RP}^2)$, hence is diffeomorphic to the entire this space (and these diffeomorphisms can be performed uniformly over all fibers). □

Lemma 17. *For arbitrary k, the group $\overline{H}_i(\sigma_9 \setminus \sigma_8, \mathbb{Q})$ is isomorphic to $\overline{H}_{i-k-11}(\mathbb{RP}^2, \mathbb{Q} \otimes \mathbb{O}r)$, i.e. to $\mathbb{Q}$ if $i = k + 13$ and to the trivial group in all other dimensions.*

Proof. The space $\sigma_9 \setminus \sigma_8$ is fibered over J_9; its fiber over a conic $C \in J_9$ is equal to $\breve{\Psi}(C) \times L(C)$, $L(C) \simeq \mathbb{R}^k$.

The spaces $L(C)$ form a trivial vector bundle over J_9. Indeed, any of its one-dimensional summands can be canonically oriented, because the space of quadratic forms vanishing on a fixed non-empty cone is oriented in an invariant way (say, from quadratic forms of signature $(2, 1)$ to those of signature $(1, 2)$).

By Proposition 2, the link $\partial\Psi(C)$ is homeomorphic to S^7. In particular, the Borel–Moore homology groups of the fibers $\breve{\Psi}(C)$ are isomorphic to $\mathbb{Q}$ in dimension 8, and are trivial in all other dimensions. By Lemma 7 the fundamental group of the base J_9 acts trivially on these homology groups.

Now, our lemma follows from the Thom isomorphism $\overline{H}_i(\sigma_9 \setminus \sigma_8, \mathbb{Q}) \simeq \overline{H}_{i-k-8}(J_9, \mathbb{Q})$ of our orientable fiber bundle over the space of conics, and the Thom isomorphism $\overline{H}_j(J_9, \mathbb{Q}) \simeq H_{j-3}(\mathbb{RP}^2, \mathbb{Q} \otimes \mathbb{O}r)$ for the fibration of J_9 over $\mathbb{RP}^2$, see Lemma 16. □

This proves the columns $p = 8$ of both tables in Fig. 1. In particular, we see that in the case of odd k both groups $E^1_{8,5} \equiv \overline{H}_{k+13}(\sigma_9 \setminus \sigma_8, \mathbb{Q})$ and $E^1_{7,5} \equiv \overline{H}_{k+12}(\sigma_8 \setminus \sigma_7, \mathbb{Q})$ are equal to $\mathbb{Q}$.



FIG. 3. Different approximations of J_7 from J_8

4.9. **The differential $d_1 : E^1_{8,k+5} \to E^1_{7,k+5}$ for odd k.**

Proposition 4. *The boundary homomorphism*

$$\overline{H}_{k+13}(\sigma_9 \setminus \sigma_8, \mathbb{Q}) \xrightarrow{\partial} \overline{H}_{k+12}(\sigma_8 \setminus \sigma_7, \mathbb{Q})$$

from the exact sequence of the triple is an isomorphism for any odd k.

Proof. Let X be an arbitrary point of the stratum J_8, i.e. a pair of crossing lines in $\mathbb{RP}^2$. This stratum has codimension 1 in the space of conics, and the open stratum J_9 approaches it at X from two different sides, see Fig. 3. Recall that $\sigma_9 \setminus \sigma_8$ is fibered over J_9 with the fiber over $C \in J_9$ equal to $\breve{\Psi}(C) \times L(C)$, $L(C) \simeq \mathbb{R}^k$. The base J_9 is oriented as an open subvariety of the space $\mathbb{RP}^5$ of all conics, the canonical orientation of the fibers $L(C)$ (and moreover of any of their k summands) was specified in the proof of Lemma 17, and the canonical orientation of fibers $\breve{\Psi}(C)$ was defined in the proof of Lemma 7. In a neighborhood U of the point X, the manifold J_8 can be considered as the boundary of any of two local components, into which it separates J_9; therefore the restriction of the orientation of J_9 to any of these components induces the boundary orientation on $J_8 \cap U$.

Lemma 18. *When the points $C \in J_9$ tend from two different local components of J_9 to a point X of the hypersurface $J_8 \subset \mathbb{RP}^5$,*

1. Two boundary orientations of J_8 at X, induced from the restrictions of one and the same orientation of J_9 to these components, are opposite to one another;

2. The 7-dimensional fundamental classes of canonically oriented fibers $\partial\Psi(C) \sim S^7$ tend to the fundamental class of the fiber $\partial\Psi(X) \setminus \Psi_6$ supplied with the canonical orientation (specified in the last paragraph of the proof of Lemma 14) and generating the Borel–Moore homology group of entire $\partial\Psi(X)$.

3. The canonical orientations of any of k summands of the k-dimensional vector bundle with fibers $L(C)$ tend to opposite orientations of the similar line over X.

By the boundary isomorphisms, item 2 of Lemma 18 is equivalent to the following statement: the 8-dimensional fundamental classes of canonically oriented fibers $\breve{\Psi}(C)$ (homeomorphic to open 8-dimensional balls) tend to the canonically oriented fundamental class of the piece of $\breve{\Psi}(X)$ generating its 8-dimensional homology group.

Proof of Lemma 18. 1. Statement 1 follows from the fact that the union of J_9 and J_8 is an open subset in the orientable manifold $\mathbb{RP}^5$ of all conics.

2. These orientations, as they were defined, depend only on the directions of the shifts of the points of 4-configurations. For J_9, it are the directions following one and the same orientation of our conic, for J_8 they go towards the crossing point, see Fig. 3. The limit positions of the former directions, when conics degenerate to a cross, differ from the latter ones an even number of times, therefore the corresponding orientations do coincide.

3. For any point $C \in J_9$ any such 1-dimensional summand consists of all quadratic functions vanishing on the corresponding cone. The basic vector, defining the canonical orientation of this line, is a quadratic function positive inside the simply-connected component of the complement of the cone in $\mathbb{R}^3$. The limits of such functions for the left- and right-hand parts of Fig. 3 are functions vanishing on the coordinate cross, but having opposite signs outside it; therefore they define opposite orientations of the corresponding summand of the fiber $L(X)$ over the point $X \in J_8$. □

So, in total the boundary orientations of $\sigma_8 \setminus \sigma_7$, induced from these two sides, differ by the factor $(-1) \times 1 \times (-1)^k$, which in the case of odd k is equal to 1 and proves the non-triviality of the boundary operator, i.e. the assertion of Proposition 4. □

4.10. **Ninth and last column.**

Proposition 5. *The group $\overline{H}_i(\sigma_{10} \setminus \sigma_9, \mathbb{Q})$ is equal to $\mathbb{Q}$ if $i = 14$ and is trivial in all other dimensions.*

Proof. The space J_{10} consists of unique point $[\mathbb{RP}^2]$ specified by the zero quadratic form. Therefore $\sigma_{10} \setminus \sigma_9 = \breve{\Psi}([\mathbb{RP}^2])$. The order complex $\Psi([\mathbb{RP}^2])$ can be calculated as the space σ of our simplicial resolution



FIG. 4. E^1 for $k = 0$ and $k = 1$

for the case $k = 0$ (i.e., with 0-dimensional spaces $L(X)$). The Borel–Moore homology group of its link $\partial\Psi([\mathbb{RP}^2]) \subset \sigma_9$ can thus be calculated by the spectral sequence, whose term E_1 coincides with columns $p \in [1, 8]$ of our main spectral sequence for $k = 0$. These columns are calculated in the previous sections, and the spectral sequence looks as is shown in the left-hand part of Fig. 4.

Lemma 19. *The differential $\partial_2 : E^2_{6,3} \to E^2_{4,4}$ of the spectral sequence of Fig. 4 (left) is an isomorphism.*

Proof. Consider our main spectral sequence for $k = 1$. All its columns with $p = 1, \dots, 8$ are already calculated, therefore it looks as is shown in the right-hand part of Fig. 4, where both signs ?? in the ninth column should be replaced by $\mathbb{Q}$ or 0 depending on whether the differential in question is trivial or not. However, the answer in the case $k = 1$ is known: the space $W^1 \setminus \Sigma$ consists of two contractible components. Therefore by Alexander duality the group $\overline{H}_i(\Sigma, \mathbb{Q}) \equiv \overline{H}_i(\sigma, \mathbb{Q})$ is equal to $\mathbb{Q}$ for $i = 5$ and is trivial in all other dimensions. Thus all cells of our spectral sequence except for $E_{1,4}$ should be killed by some its differential. This is impossible if the group $E^1_{9,0}$ is non-trivial. □

So, $H_i\big(\partial\Psi([\mathbb{RP}^2]), \mathbb{Q}\big)$ is equal to $\mathbb{Q}$ for $i = 0$ and $i = 13$, and is trivial in all other dimensions. Proposition 5 follows from this fact by the exact sequence of the pair $\big(\Psi([\mathbb{RP}^2]), \partial\Psi([\mathbb{RP}^2])\big)$. □

Theorem 2 is completely proved. □

5. Concluding remarks

1. A problem: is there a more direct proof of Theorem 1, in the same way as Proposition 1 can be proved by the methods of [5] type?

Of course, the space $W^k \setminus \Sigma$ is fibered over S^{k-1}: with any non-resultant system we can associate the image of a fixed vector, say $(1,0,0)$. However, it seems that this structure helps us quite little. The Poincaré polynomials of the fibers of these fiber bundles follow easily from Theorem 1: in the case of odd (respectively, even) $k \geq 3$ they are equal to $1+t^{3k-7}(1+t^{k-5})(1+t^{k-2})$ (respectively, $1+t^{3k-9}+t^{5k-14}$).

2. The construction of resolutions of discriminants in the terms of self-joins was formulated in full generality (sufficient for our current needs) in [10] (although its particular versions appeared in less difficult situations also in [16], [17], [19]). It may seem that this construction is less canonical than the one in terms of Hilbert schemes (see e.g. [20]). However in practice, when one needs not only to prove abstract theorems, but to accomplish a difficult explicit homological calculation, the approach of [10] proves to be more effective.

3. The formal analog. The space W^k of quadratic maps $\mathbb{R}^3 \to \mathbb{R}^k$ is a subspace of the (infinite-dimensional) space of all maps $\mathbb{R}^3 \to \mathbb{R}^k$, quadratic in restriction to any one-dimensional subspace in $\mathbb{R}^3$. The latter space of maps also contains a resultant variety: it consists of maps having common zeros in $\mathbb{R}^3 \setminus 0$. The corresponding space of non-resultant maps is homotopy equivalent to the space of continuous maps $\mathbb{RP}^2 \to S^{k-1}$. Its rational homology group for $k > 3$ can be easily calculated (e.g. by the methods of [17]) and coincides with that of S^{k-1}: all other rational homology groups are trivial by Corollary 1 of the present article. So it is much poorer than its nonstable analog calculated above.

References

[1] V.I. Arnold, *On some topological invariants of algebraic functions,* Transact. Moscow Math. Soc. **21** (1970), p. 30–52.

[2] V.I. Arnold, *Spaces of functions with moderate singularities,* Funct. Anal. and its Appl. **23**:3, (1989), p. 1–10.

[3] A. Bahri, J.M. Coron, *On a nonlinear elliptic equation involvingthe critical Sobolel exponent: the effect of the topology of the domain*, Comm. Pure and Applied Mathematics, **XLI** (1988), 253–294.

[4] L. Blum, M. Shub, S. Smale, *On a theory of computations over the real numbers: NP completeness, recursive functions, and universal machines.* Bull. Amer. Math. Soc., **21**(1) (1989), 1–46.

[5] A. Borel, *Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts,* Ann. Math., **57** (1953), 115–207.

[6] A. Borel, J.-P. Serre, *Cohomologie d'immeubles et de groupes S-arithmetiques*, Topology **15** (1978), 211–231.

[7] F.R. Cohen, R.L. Cohen, B.M. Mann, R.J. Milgram, *The topology of the space of rational functions and divisors of surfaces,* Acta Math., **166** (1991), 163–221.

[8] S. Kallel, R. Karoui, *Symmetric joins and weighted barycenters*, Advanced Nonlinear Studies **11** (2011), 117–143; arXiv:math/0602283v3

[9] A. Kozlowski and R. Yamaguchi, *Topology of complements of discriminants and resultants,* Journal of the Mathematical Society of Japan, **52**:4 (2000), 949-959.

[10] A. Gorinov, *Real cohomology groups of the space of nonsingular curves of degree 5 in* $\mathbb{CP}^2$*,* Ann. Fac. Toulouse Math. (6), 14(3) (2005), 395–434. arXiv: math.AT/0105108

[11] J.-P. Serre, *Homologie singulìere des espaces fibrés. Applications,* Ann. Math., **54** (1951), 425–505.

[12] M. Shub, S. Smale, *On the intractability of Hilbert's Nullstellensatz and an algebraic version of "*$NP \neq P$*?"*, Duke Math. J., **81**:1 (1995), 47–54.

[13] S. Smale, *Mathematical Problems for the next Century,* Mathematical Intelligencer **20**(2), 7-15.

[14] O. Tommasi, *Rational cohomology of the moduli space of genus 4 curves*, Compos. Math., **141**(2) (2005), 359-384.

[15] O. Tommasi, *Geometry of Discriminants and Cohomology of Moduli Spaces*, Thesis, Radboud Universiteit Nijmegen, 2005. http://repository.ubn.ru.nl/bitstream/handle/2066/27027/27027.pdf?sequence=1

[16] V.A. Vassiliev, *Stable cohomology of complements to the discriminants of deformations of singularities of smooth functions*, Itogi Nauki i Tekhn. Ser. Sovrem. Probl. Math., Nov. Dostizh., **33** (1988), 3–29, English translation in: J. of Soviet Math., **52:4** (1990), 3217–3230.

[17] V.A. Vassiliev, *Complements of discriminants of smooth maps: topology and applications. Revised ed.*, Transl. of Math. Monographs, v. 98, AMS, Providence RI., 1994.

[18] V.A. Vassiliev, *Topology of complements of discriminants,* Moscow, Phasis, 1997, 552 p. (in Russian).

[19] V.A. Vassiliev, *Homology of spaces of homogeneous polynomials in* $\mathbb{R}^2$ *without multiple zeros*, Proc. Steklov Mathematical Institute, **221** (1998), 133–138; arXiv: 1407.7230.

[20] V.A. Vassiliev, *How to calculate the homology of spaces of nonsingular algebraic projective hypersurfaces*, Proc. Steklov Mathematical Institute, **225** (1999), 121–140; arXiv: 1407.7229.

[21] V.A. Vassiliev, *Homology of spaces of non-resultant polynomial systems in* $\mathbb{R}^2$ *and* $\mathbb{C}^2$*,* Arnold Mathematical Journal, **1**:3 (2015), 233–242, arXiv: 1409.6005.

[22] V.A. Vassiliev, *Rational homology of the order complex of zero sets of homogeneous quadratic polynomial systems in* $\mathbb{R}^3$, Proc. Steklov Inst. Math., **290** (2015), 197–209; arXiv: 1501.06063.

Steklov Mathematical Institute of Russian Academy of Sciences; National Research Institute — Higher School of Economics

E-mail address: vva@mi.ras.ru